

RESEARCH NOTE

Clinical epidemiology of ciprofloxacin-resistant *Proteus mirabilis* isolated from urine samples of hospitalised patients

R. Saito¹, S. Okugawa¹, W. Kumita², K. Sato²,
T. Chida², N. Okamura², K. Moriya¹ and
K. Koike¹

¹Department of Infection Control and Prevention,
The University of Tokyo Hospital and

²Department of Microbiology and Immunology,
Graduate School of Health Sciences, Tokyo
Medical and Dental University, Tokyo, Japan

ABSTRACT

This study investigated the clinical characteristics of ciprofloxacin-resistant *Proteus mirabilis* isolates from urine samples associated with nosocomial infection or colonisation, and identified the risk-factors for ciprofloxacin resistance. Data for patients with ciprofloxacin-resistant *P. mirabilis* isolates ($n = 13$) were compared with those for randomly selected patients with ciprofloxacin-susceptible *P. mirabilis* isolates ($n = 40$) who were matched by temporal occurrence as control patients. The majority of ciprofloxacin-resistant *P. mirabilis* isolates were multiresistant, and ciprofloxacin resistance was associated significantly with previous use of fluoroquinolones and production of extended-spectrum β -lactamases.

Keywords Epidemiology, extended-spectrum β -lactamases, fluoroquinolones, *Proteus mirabilis*, risk-factors, urinary tract infection

Original Submission: 13 April 2007; **Revised Submission:** 22 June 2007; **Accepted:** 11 July 2007

Clin Microbiol Infect 2007; **13**: 1204–1206
10.1111/j.1469-0691.2007.01826.x

Fluoroquinolones are potent antimicrobial agents that are used in the treatment of a wide

range of community-acquired and nosocomial infections. However, resistance to fluoroquinolones has increased significantly since the late 1980s [1], and it has been demonstrated that use of fluoroquinolones is an important risk-factor for ciprofloxacin resistance in *Escherichia coli* and *Klebsiella pneumoniae* [2,3]. In addition, following the worldwide use of broad-spectrum cephalosporins during the past two decades, Gram-negative bacteria producing extended-spectrum β -lactamases (ESBLs) have emerged. This emerging resistance is of great concern because its association with resistance to other unrelated antimicrobial agents severely limits therapeutic options [2].

Proteus mirabilis is one of the most common causes of urinary tract infection, and infections caused by this organism are often nosocomial, persistent and difficult to treat. Although wild-type strains of *P. mirabilis* are usually susceptible to fluoroquinolones and β -lactams, a progressive increase in resistance to fluoroquinolones and broad-spectrum cephalosporins has been seen in clinical isolates of this species [4–6]. The relationship between fluoroquinolone resistance and ESBL production in *E. coli* and *K. pneumoniae* is now well-known [3,7,8], but the epidemiology of fluoroquinolone resistance and its relationship with ESBL production in *P. mirabilis* has not yet been clarified. The present study therefore investigated the clinical characteristics of ciprofloxacin-resistant *P. mirabilis* isolates from urine and identified the risk-factors for ciprofloxacin resistance.

The study was conducted at a 1193-bed teaching hospital, providing care for up to 1 million individuals, between April 2003 and March 2006. The medical records of patients were reviewed retrospectively, and all patients with *P. mirabilis* isolated from urine were enrolled in the study. An infectious disease physician evaluated all patients and classified the *P. mirabilis* isolates according to CDC criteria for nosocomial infection [9]. Colonisation was defined when these criteria were not fulfilled. Analysis results for consecutive urine samples containing $\geq 10^5$ CFU/mL with pyuria (≥ 10 leukocytes/high-power field) were obtained from medical records. Data were also obtained for patients with ciprofloxacin-resistant *P. mirabilis* isolates and for randomly selected patients with ciprofloxacin-susceptible *P. mirabilis* isolates, who were matched by temporal occurrence as control

Corresponding author and reprint requests: R. Saito, Department of Infection Control and Prevention, The University of Tokyo Hospital, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan
E-mail: saito-lab@h.u-tokyo.ac.jp

patients. Data recorded were age, gender, underlying disease, date of hospitalisation, presence of a urinary catheter or a central venous catheter, and antibiotic therapy within the preceding 90 days.

All isolates were identified using the Vitek I system (bioMérieux, Tokyo, Japan). Susceptibility to ciprofloxacin, levofloxacin, minocycline, ampicillin, ampicillin-sulbactam, piperacillin, ceftazidime, cefotaxime, aztreonam, amikacin, gentamicin, imipenem and meropenem was tested using panels manufactured by Eiken Chemical (Tokyo, Japan). Nalidixic acid (Wako Pure Chemical Industries, Osaka, Japan) was also used. MICs were determined by broth micro-dilution [10]. Quality control organisms for susceptibility tests were *Staphylococcus aureus* ATCC 21293, *E. coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853. ESBL production was initially detected using cefotaxime and cefotaxime-clavulanate disks, with ESBL-producing *P. mirabilis* isolates being further screened for the presence of class A β -lactamase genes (*bla*_{TEM}, *bla*_{SHV} and *bla*_{CTX-M}) by PCR using the primer sets described previously [11,12]. Purified PCR fragments were sequenced using an ABI Prism 310 DNA sequencer (Applied Biosystems, Foster City, CA, USA), followed by a similarity search using the BLAST program (DDBJ, Shizuoka, Japan). Clonal relationships among ciprofloxacin-resistant *P. mirabilis* isolates were investigated by pulsed-field gel electrophoresis following digestion of chromosomal DNA using *NotI* (Roche Diagnostics, Tokyo, Japan). Data were analysed using JMP software, v.6.0 (SAS Institute, Tokyo, Japan) and categorical variables were compared by means of either the chi-square test or Fisher's exact test, as appropriate. A two-tailed p value of <0.05 was considered to be significant.

Among the 80 non-repetitive *P. mirabilis* isolates from urine samples, 13 (16%) were ciprofloxacin-resistant. Of the 13 patients involved, three (23%) were classified as being infected, compared with seven (18%) of 40 control patients (not significant). The antibiotic susceptibilities of the *P. mirabilis* isolates are shown in Table 1. The ciprofloxacin-resistant isolates showed a significantly higher frequency of resistance to ampicillin, piperacillin, ampicillin-sulbactam, cefotaxime, nalidixic acid and levofloxacin ($p < 0.001$). Univariate analysis revealed that previous use of fluoroquinolones ($p 0.042$) and

Table 1. Antibiotic susceptibilities and risk-factors for *Proteus mirabilis* isolates from 13 patients (cases) with ciprofloxacin-resistant isolates and 40 patients (controls) with ciprofloxacin-susceptible isolates

	Cases (<i>n</i> = 13)	Controls (<i>n</i> = 40)	<i>p</i>
Antibiotic resistance frequency, <i>n</i> (%)			
Ampicillin	11 (85)	1 (3)	<0.001
Piperacillin	8 (62)	0	<0.001
Ampicillin-sulbactam	11 (85)	0	<0.001
Ceftazidime	0	0	NS
Cefotaxime	8 (62)	0	<0.001
Aztreonam	1 (8)	0	NS
Gentamicin	0	0	NS
Amikacin	0	0	NS
Imipenem	0	0	NS
Meropenem	0	0	NS
Minocycline	11 (85)	25 (63)	NS
Nalidixic acid	13 (100)	9 (23)	<0.001
Levofloxacin	8 (62)	0	<0.001
Risk-factors, <i>n</i> (%)			
Age, median years (range)	73 (53–91)	65 (3–94)	NS
Male	7 (54)	14 (35)	NS
Diabetes mellitus	3 (23)	9 (23)	NS
Urinary catheter	6 (46)	18 (45)	NS
Central venous catheter	3 (23)	5 (13)	NS
Long hospital stay (>2 weeks)	5 (38)	14 (35)	NS
ESBL production	8 (62)	0	<0.001
Use of other antibiotics within the previous 90 days	7 (54)	12 (30)	NS
Cephems	1 (8)	8 (20)	NS
Penicillins	2 (15)	1 (3)	NS
Aminoglycosides	1 (8)	1 (3)	NS
Fluoroquinolones	3 (23)	1 (3)	0.042
Carbapenems	2 (15)	2 (5)	NS

ESBL, extended-spectrum β -lactamase; NS, not significant.

ESBL production ($p < 0.001$) were associated significantly with ciprofloxacin resistance (Table 1). Eight isolates produced a CTX-M-2 ESBL. Eleven genotypes were identified by pulsed-field gel electrophoresis among the 13 ciprofloxacin-resistant *P. mirabilis* isolates, with three isolates being clonally related.

Infections caused by fluoroquinolone-resistant ESBL-producing *E. coli* and *K. pneumoniae* strains have now been reported throughout the world. It has also been demonstrated that 18–56% of ESBL-producing *E. coli* and *K. pneumoniae* strains are resistant to fluoroquinolones, and that fluoroquinolone resistance is associated closely with ESBL production [3,7,8]. Another cause for concern is a marked increase in the incidence of *P. mirabilis* isolates that are resistant to fluoroquinolones and broad-spectrum cephalosporins [4–6]. Previous studies have demonstrated that ESBL-producing isolates are usually resistant to penicillins, cephalosporins, aminoglycosides and trimethoprim-sulphamethoxazole [3,6]. The present study revealed that the majority of ciprofloxacin-resistant *P. mirabilis* isolates from urine samples were multiresistant, with most being resistant to cefotaxime and levofloxacin, but not to aminoglycosides.

In the present study, acquisition of ciprofloxacin resistance by *P. mirabilis* was found to be associated significantly with previous use of fluoroquinolones and ESBL production, as was also found for ESBL-producing *E. coli* and *K. pneumoniae* [3,7,8]. While the basis for combined ciprofloxacin and broad-spectrum cephalosporin resistance is not yet fully understood, the presence of both ciprofloxacin resistance and ESBL production may be a consequence of the interplay of previous heavy antibiotic use and conditions that favour patient-to-patient transfer of multidrug-resistant organisms [3]. Other possible explanations include active efflux, outer-membrane alterations and ciprofloxacin resistance mediated by plasmids containing a class A β -lactamase gene.

In conclusion, the present study has certain limitations, in that it was retrospective and sample numbers were small. However, the ciprofloxacin-resistant *P. mirabilis* isolates also exhibited multiresistance, and an analysis of risk-factors showed that previous use of fluoroquinolones and ESBL production were the strongest determinants of the acquisition of ciprofloxacin resistance by *P. mirabilis*. Therefore, when treatment protocols are designed for patients following the isolation of ciprofloxacin-resistant *P. mirabilis* from urine samples, the prevalence of ESBL-producing strains should be taken into consideration, and empirical antibiotic treatments with greater efficacy should be selected accordingly.

REFERENCES

1. Richard P, Delangle MH, Raffi F *et al.* Impact of fluoroquinolone administration on the emergence of fluoroquinolone-resistant gram-negative bacilli from gastrointestinal flora. *Clin Infect Dis* 2001; **32**: 162–166.
2. Hyle EP, Lipworth AD, Zaoutis TE *et al.* Risk factors for increasing multidrug resistance among extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella* species. *Clin Infect Dis* 2005; **40**: 1317–1324.
3. Paterson DL, Mulazimoglu L, Casellas JM *et al.* Epidemiology of ciprofloxacin resistance and its relationship to extended-spectrum beta-lactamase production in *Klebsiella pneumoniae* isolates causing bacteremia. *Clin Infect Dis* 2000; **30**: 473–478.
4. Hernandez JR, Martinez-Martinez L, Pascual A *et al.* Trends in the susceptibilities of *Proteus mirabilis* isolates to quinolones. *J Antimicrob Chemother* 2000; **45**: 407–408.
5. Endimiani A, Luzzaro F, Brigante G *et al.* *Proteus mirabilis* bloodstream infections: risk factors and treatment outcome related to the expression of extended-spectrum beta-lactamases. *Antimicrob Agents Chemother* 2005; **49**: 2598–2605.
6. Kim JY, Park YJ, Kim SI *et al.* Nosocomial outbreak by *Proteus mirabilis* producing extended-spectrum beta-lactamase VEB-1 in a Korean university hospital. *J Antimicrob Chemother* 2004; **54**: 1144–1147.
7. Lautenbach E, Strom BL, Bilker WB *et al.* Epidemiological investigation of fluoroquinolone resistance in infections due to extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. *Clin Infect Dis* 2001; **33**: 1288–1294.
8. Tolun V, Kucukbasmaci O, Torumkuney-Akbulut D *et al.* Relationship between ciprofloxacin resistance and extended-spectrum beta-lactamase production in *Escherichia coli* and *Klebsiella pneumoniae* strains. *Clin Microbiol Infect* 2004; **10**: 72–75.
9. Garner JS, Jarvis WR, Emori TG *et al.* CDC definitions for nosocomial infections. *Am J Infect Control* 1988; **16**: 128–140.
10. Clinical and Laboratory Standards Institute. *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically*, approved standard M7-A7. Wayne, PA: CLSI, 2006.
11. Yagi T, Kurokawa H, Shibata N *et al.* A preliminary survey of extended-spectrum β -lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* and *Escherichia coli* in Japan. *FEMS Microbiol Lett* 2000; **184**: 53–56.
12. Pitout JD, Hossain A, Hanson ND. Phenotypic and molecular detection of CTX-M- β -lactamases produced by *Escherichia coli* and *Klebsiella* spp. *J Clin Microbiol* 2004; **42**: 5715–5721.